arXiv:1202.2398v1 [math.FA] 11 Feb 2012

THE POMPEIU PROBLEM AND DISCRETE GROUPS

MICHAEL J. PULS

Abstract. We formulate a version of the Pompeiu problem in the discrete group setting. Necessary and sufficient conditions are given for a finite collection of finite subsets of a discrete abelian group, whose torsion free rank is less than the cardinal of the continuum, to have the Pompeiu property. We also prove a similar result for nonabelian free groups.

1. Introduction

Let $\mathbb{C}$ be the complex numbers, $\mathbb{R}$ the real numbers, $\mathbb{Z}$ the integers and $\mathbb{N}$ the natural numbers. Let $\mathcal{K}$ be a finite family of compact subsets of $\mathbb{R}^2$. Denote by $\mathcal{T}$ the group of translations on $\mathbb{R}^2$. One version of the Pompeiu problem asks under what conditions on $\mathcal{K}$ will $f = 0$ be the only continuous function on $\mathbb{R}^2$ that satisfies

$$\int_{\tau(K)} f\, dx = 0$$

for all $K \in \mathcal{K}$ and for all $\tau \in \mathcal{T}$. Variations of the Pompeiu problem have been studied in various contexts, see [1, 3, 9, 11, 12, 13] and the references therein for more information. The purpose of this study is to investigate a version of the Pompeiu problem in the discrete group setting that is similar to the above problem.

For the rest of this paper G will always denote a discrete group. So, the compact subsets of G are precisely the finite subsets of G and all complex-valued functions on G are continuous. Let $\mathcal{K}$ be a finite collection of finite subsets of G. We now state the Pompeiu problem for discrete groups with respect to left translations.

Problem 1.1. *When is $f = 0$ the only complex-valued function from a given class of functions on G that satisfies*

$$\sum_{x \in gK} f(x) = 0 \tag{1.1}$$

for all $K \in \mathcal{K}$ and for all $g \in G$?

We shall say that $\mathcal{K}$ has the *Pompeiu property* with respect to left translations if $f = 0$ is the only function from the class of functions under consideration that

---

Date: February 10, 2012.

2000 *Mathematics Subject Classification.* Primary: 43A15; Secondary: 43A32, 43A70, 43A80, 43A90.

Key words and phrases. abelian group, free group, left translates, Pompeiu property, radial functions, radial translates, torsion free rank, zero divisors.

satisfies (1.1). For brevity, we will say that $\mathcal{K}$ has the Pompeiu property when it has the Pompeiu property with respect to left translations.

In Section 2 we give needed definitions and prove some preliminary results. In particular we make a connection between the problem of zero divisors on torsion free discrete groups and Problem 1.1. We will give some examples of groups for which (1.1) is true when we assume an extra condition on the decay of f. The torsion free rank of G, which we shall denote by $r_0(G)$, is the cardinality of a maximal independent system of elements in G of infinite order. Indicate the cardinal of the continuum by 2^ω. In Section 3 we give necessary and sufficient conditions on a finite collection of finite subsets of G to have the Pompeiu property when G is abelian and $r_0(G) < 2^\omega$. Let F_k be the free group on k generators, where $k \geq 2$. In Section 4 we give necessary and sufficient conditions on a finite set of finite radial subsets, to be defined in Section 4, of F_k to have the Pompeiu property.

2. Preliminaries and the problem of zero divisors

For $z \in \mathbb{C}$, $\bar{z}$ will denote the complex conjugate of z. We represent a complex-valued function f on G by a formal sum $f = \sum_{g\in G} a_g g$ where $a_g \in \mathbb{C}$ and $f(g) = a_g$. Also, we shall write $\tilde{f}$ for $\sum_{g\in G} a_g g^{-1}$ and $\bar{f}$ for $\sum_{g\in G} \overline{a_g} g$. Indicate by $\mathcal{F}(G)$ the set of all functions on G. For $1 \leq p \in \mathbb{R}$, $L^p(G)$ will consist of those formal sums for which $\sum_{g\in G} |a_g|^p < \infty$, and $\mathbb{C}G$, the group ring of G over $\mathbb{C}$, will consist of all formal sums that satisfy $a_g = 0$ for all but finitely many g. The group ring $\mathbb{C}G$ can also be thought of as the set of all functions on G with compact support. For $g \in G$ and $f \in \mathcal{F}(G)$, the left translate $L_g(f)$ of f is defined to be $(L_g(f))(x) = f(gx)$.

Let $\alpha = \sum_{g\in G} a_g g \in \mathbb{C}G$ and $f = \sum_{g\in G} b_g g \in \mathcal{F}(G)$. Define a map $\langle \cdot, \cdot \rangle \colon \mathbb{C}G \times \mathcal{F} \mapsto \mathbb{C}$ by

$$\langle \alpha, f \rangle = \sum_{g\in G} a_g \overline{b_g}.$$

For a fixed $h \in \mathcal{F}(G)$, $\langle \cdot, h \rangle$ is a linear functional on $\mathbb{C}G$. Now let T be a linear functional on $\mathbb{C}G$. Define $f(g) = \overline{T(g)}$ for each $g \in G$. Thus each linear functional on $\mathbb{C}G$ defines an element of $\mathcal{F}(G)$. Hence, the dual of $\mathbb{C}G$ can be identified with $\mathcal{F}(G)$. If E is a subset of G, denote by χ_E the characteristic function on E. Let $g \in G$. Observe that $L_g(\chi_E) = \chi_{g^{-1}E}$. Suppose $\sum_{x\in gK} f(x) = 0$ for all $g \in G$ and for some finite subset K of G. Then

$$\sum_{x\in gK} f(x) = \sum_{x\in G} \chi_{gK}(x) f(x) = \sum_{x\in G} (L_{g^{-1}}(\chi_K))(x) f(x) = \langle (L_{g^{-1}}(\chi_K)), \bar{f} \rangle.$$

We have just proved

Lemma 2.1. *Let K be a finite subset of G, and let f be a function on G. Then $\sum_{x\in gK} f(x) = 0$ for all $g \in G$ if and only if $\langle L_g(\chi_K), \bar{f} \rangle = 0$ for all $g \in G$.*

Lemma 2.1 will eventually allow us to give a link between Pompeiu's problem and the problem of zero divisors.

Let X represent either $\mathcal{F}(G), L^p(G)$ or $\mathbb{C}G$. For $\alpha = \sum_{g\in G} a_g g \in \mathbb{C}G$ and $f = \sum_{g\in G} b_g g \in X$, we define a multiplication, known as convolution, $\mathbb{C}G \times X \mapsto X$

by

$$\alpha * f = \sum_{g,h\in G} a_g b_h gh = \sum_{g\in G}\left(\sum_{h\in G} a_{gh^{-1}} b_h\right) g.$$

Lemma 2.2. *Let $\alpha \in \mathbb{C}G$ and let $f \in \mathcal{F}(G)$. Then $\alpha * \tilde{f} = 0$ if and only if $\langle L_g(\alpha), \bar{f}\rangle = 0$ for all $g \in G$.*

Proof. Set $\alpha = \sum_{g\in G} a_g g$ and $f = \sum_{g\in G} b_g g$. Note that for $g \in G, L_g(\alpha) = \sum_{x\in G} a_{gx} x$. Then

$$\alpha * \tilde{f} = \sum_{g\in G}\left(\sum_{x\in G} a_{gx} b_x\right) g$$

and $\langle L_g(\alpha), \bar{f}\rangle = \sum_{x\in G} a_{gx} b_x$. The result now follows. ☐

By combining Lemma 2.1 and Lemma 2.2 we obtain

Proposition 2.3. *Let $\mathcal{K}$ be a finite collection of finite subsets of G. Then $f = 0$ is the only function on G that satisfies (1.1) if and only if $f = 0$ is the only solution to the system of convolution equations*

$$\chi_K * \tilde{f} = 0 \text{ for all } K \in \mathcal{K}.$$

We just saw that there is a connection between Pompeiu's problem on discrete groups and the following problem concerning zero divisors.

Problem 2.4. *Let G be a torsion free group and let X be one of the following: $\mathbb{C}G, L^p(G)$ or $\mathcal{F}(G)$. If $0 \neq \alpha \in \mathbb{C}G$ and if $0 \neq f \in X$, then is $\alpha * f = 0$?*

We point out that if G is not torsion free, then there are zero divisors in $\mathbb{C}G$. For example, let 1 be the identity element of G and let $g \in G$ such that $g \neq 1$ and $g^n = 1$ for some $n \in \mathbb{N}$. Then $(1 + g + \cdots + g^{n-1}) * (1 - g) = 0$. It now follows that the set $K = \{1, g, \ldots, g^{n-1}\}$ does not have the Pompeiu property. To see this, set $f = 1 - g \in \mathbb{C}G$ and note $\chi_K = 1 + g + \cdots + g^{n-1}$. Using the fact $g^{-k} = g^{n-k}$ for $0 \leq k \leq n$, we have $\chi_K * \tilde{f} = 0$. By Proposition 2.3, $f = 1 - g$ satisfies (1.1) when $K = \{1, g, \ldots, g^{n-1}\}$ and $g^n = 1$.

The zero divisor problem has been studied in [2, 6, 7, 8, 10]. Using results obtained in some of these papers we can give some results concerning (1.1) for specific groups G and classes of functions on G. For example, in the case $G = \mathbb{Z}^n$, where $2 \leq n \in \mathbb{N}$, if f satisfies (1.1) and $f \in L^p(\mathbb{Z}^n)$ for $p \leq \frac{2n}{n-1}$, then by [8, Theorem 2.1] $f = 0$. Let F_k denote the free group on k generators, where $k \geq 2$. The main result of [7] shows that $f = 0$ is the only element of $L^2(F_k)$ for which (1.1) is true. In Section 4 we will see that for $p > 2$, there exists a nonzero f in $L^p(F_k)$ that satisfies (1.1) for certain subsets of F_k that are circular.

3. ABELIAN GROUPS

In this section G will always be a discrete abelian group with $r_0(G) < 2^\omega$. We will give necessary and sufficient conditions for a finite set of finite subsets of G to have the Pompeiu property.

Let M be a maximal ideal in $\mathbb{C}G$. Then $\mathbb{C}G/M$ is a field since $\mathbb{C}G$ is a commutative ring with identity 1. Let $i_M\colon \mathbb{C}G \mapsto \mathbb{C}G/M$ be the natural map. It was shown in the proof of Theorem 3 in [5] that $\mathbb{C}G/M$ is isomorphic to $\mathbb{C}$. Thus i_M can be considered a ring homomorphism from $\mathbb{C}G$ onto $\mathbb{C}$ with kernel M. We define a function $f_M\colon G \to \mathbb{C}$ by

$$f_M(g) = i_M(g)$$

for a given maximal ideal M of $\mathbb{C}G$. Each $g \in G$ is invertiable in $\mathbb{C}G$. Thus $f_M(g) \neq 0$ since $i_M(g) \neq 0$ in $\mathbb{C}G/M$. For $\alpha \in \mathbb{C}G$, define

$$Z(\alpha)\colon = \{i_M \mid i_M(\alpha) = 0\},$$

where M is a maximal ideal in $\mathbb{C}G$. For an ideal I in $\mathbb{C}G$, set $Z(I) = \cap_{\alpha\in I} Z(\alpha)$.

Proposition 3.1. *Let G be a discrete abelian group with $r_0(G) < 2^\omega$. Let $\alpha = \sum_{g\in G} a_g g \in \mathbb{C}G$ and let M be a maximal ideal in $\mathbb{C}G$. Then $\alpha \in M$ if and only if $\alpha * \widetilde{f_M} = 0$.*

Proof. Suppose $\alpha \in M$. Then $i_M(\alpha) = 0$. Because $L_x(\alpha) = x^{-1} * \alpha$ for $x \in G$ and M is an ideal, $i_M \in Z(L_x(\alpha))$ for each $x \in G$. We now obtain

$$0 = i_M(L_x(\alpha)) = \sum_{g\in G} a_{xg} i_M(g) = \langle L_x(\alpha), \overline{f_M}\rangle$$

for all $x \in G$. By Lemma 2.2 $\alpha * \widetilde{f_M} = 0$.

Conversely, let $\alpha \in \mathbb{C}G$ and assume $\alpha * \widetilde{f_M} = 0$ for some maximal ideal M in $\mathbb{C}G$. Lemma 2.2 says that $\langle L_x(\alpha), \overline{f_M}\rangle = 0$ for all $x \in G$. In particular

$$\langle \alpha, \overline{f_M}\rangle = \sum_{g\in G} a_g i_M(g) = i_M(\alpha).$$

Thus, $\alpha \in M$. □

We now state and prove a Wiener type result.

Theorem 3.2. *Let G be a discrete abelian group with $r_0(G) < 2^\omega$ and let I be an ideal in $\mathbb{C}G$. Then $Z(I) = \emptyset$ if and only if $I = \mathbb{C}G$.*

Proof. Assume $Z(I) \neq \emptyset$. Then there exists a maximal ideal M in $\mathbb{C}G$ for which $i_M(\alpha) = 0$ for each $\alpha \in I$. The kernel of i_M is precisely M, so $\alpha \in M$. By Proposition 3.1, $\alpha * \widetilde{f_M} = 0$ for all $\alpha \in I$. Thus $1 \notin I$ because $1 * \widetilde{f_M} = \widetilde{f_M}$ and $\widetilde{f_M} \neq 0$. Hence, $I \neq \mathbb{C}G$.

Now suppose $I \neq \mathbb{C}G$. Let M be a maximal ideal that contains I. Then $i_M(\alpha) = 0$ for each $\alpha \in I$. Hence $i_M \in Z(I)$. □

Corollary 3.3. *Let G be a discrete abelian group with $r_0(G) < 2^\omega$ and let $\mathcal{K}$ be a finite set of finite subsets of G. Then $\mathcal{K}$ has the Pompeiu property if and only if $\cap_{K\in\mathcal{K}} Z(\chi_K) = \emptyset$.*

Proof. Assume $\cap_{K\in\mathcal{K}} Z(\chi_K) = \emptyset$. Let I be the ideal in $\mathbb{C}G$ generated by the functions $\chi_K, K \in \mathcal{K}$. Then $Z(I) = \emptyset$. Suppose that $\mathcal{K}$ does not have the Pompeiu property. Proposition 2.3 says that there exists a nonzero function $f \in \mathcal{F}(G)$ for which $\chi_K * \tilde{f} = 0$ for all $K \in \mathcal{K}$. Thus $I \neq \mathbb{C}G$. Using Theorem 3.2 we obtain the contradiction $Z(I) \neq \emptyset$. Thus $\mathcal{K}$ has the Pompeiu property.

Conversely, assume $\cap_{K\in\mathcal{K}} Z(\chi_K) \neq \emptyset$. Then there exists a maximal ideal M in $\mathbb{C}G$ such that $i_M \in Z(\chi_K)$ for each $K \in \mathcal{K}$. Thus $\chi_K \in M$ for each $K \in \mathcal{K}$. By combining Proposition 3.1 and Proposition 2.3 we obtain that $\mathcal{K}$ does not have the Pompeiu property. □

4. Free groups

Let F_k denote the free group on k generators, where $2 \leq k \in \mathbb{N}$. In this section we derive results for F_k that are analogous to those given in Section 3 for abelian groups. In particular, we give necessary and sufficient conditions on a finite collection $\mathcal{K}$ of certain subsets of F_k so that $\mathcal{K}$ will have the Pompeiu property. We conclude the section with a discussion about how our work is connected with some of the results in [3].

Indicate by e the identity element of F_k. Any element x of F_k has an unique expression as a finite product of generators and their inverses, which does not contain any two adjacent factors ww^{-1} or $w^{-1}w$. The number of factors in x is called the *length* of x and is denoted by $|x|$. The identity element e will have length zero.

A function in $\mathcal{F}(F_k)$ is said to be *radial* if its value depends only on $|x|$. Let $E_n = \{x \in F_k \mid |x| = n\}$, and let e_n indicate the cardinality of E_n. Then $e_n = 2k(2k-1)^{n-1}$ for $n \geq 1$, and $e_0 = 1$. We shall say that a subset A of F_k is a *radial set* if A is the union of sets of the form E_n. Let χ_n denote the characteristic function on E_n, so as an element of $\mathbb{C}F_k$ we have $\chi_n = \sum_{|x|=n} x$. Then every radial function has the form $\sum_{n=0}^{\infty} a_n \chi_n$, where $a_n \in \mathbb{C}$. Let $(\mathbb{C}F_k)_r$ denote the radial functions contained in $\mathbb{C}F_k$ and let $\omega = \sqrt{2k-1}$. It was shown in [4, Chapter 3] that

$$\begin{aligned} \chi_1 * \chi_1 &= \chi_2 + 2k * \chi_0 \\ \chi_1 * \chi_n &= \chi_{n+1} + \omega^2 \chi_{n-1}, \quad n \geq 2, \end{aligned}$$

hence $(\mathbb{C}F_k)_r$ is a commutative ring which is generated by χ_0 and χ_1.

A slight modification to the proof of [8, Lemma 6.1] gives the following elementary result, which will be needed later.

Lemma 4.1. *Let $x, y \in F_k$, with $|x| = |y|$, and let $0 \leq m, n \in \mathbb{Z}$. Then*

$$\langle x * \chi_m, \chi_n \rangle = \langle y * \chi_m, \chi_n \rangle.$$

For $0 \leq n \in \mathbb{Z}$ define polynomials $P_n(z)$ by

$$P_0(z) = 1, \quad P_1(z) = z, \quad P_2(z) = z^2 - 2k$$
$$\text{and } P_{n+1}(z) = zP_n(z) - \omega^2 P_{n-1}(z) \text{ for } n \geq 2.$$

Using the above relations with an induction argument it follows

$$\chi_n = P_n(\chi_1).$$

For $\alpha = \sum_{j=0}^{m} a_j \chi_j \in (\mathbb{C}F_k)_r$, define a function $\hat{\alpha}\colon \mathbb{C} \to \mathbb{C}$ by

$$\hat{\alpha}(z) = \sum_{j=0}^{m} a_j P_j(z).$$

Let $Z(\alpha) = \{z \mid \hat{\alpha}(z) = 0\}$. If I is an ideal in $(\mathbb{C}F_k)_r$, then set $Z(I) = \cap_{\alpha \in I} Z(\alpha)$.

A function ψ on F_k is said to be *spherical* if and only if it defines a multiplicative linear functional on $(\mathbb{C}F_k)_r$. See pages 33-34, especially Lemma 1.5, of [4] for a more detailed discussion about spherical functions on F_k. Let $z \in \mathbb{C}$ and define $\phi_z \in \mathcal{F}(F_k)$ by

$$\phi_z = \sum_{n=0}^{\infty} \frac{P_n(z)}{e_n} \chi_n.$$

Now,

$$\begin{aligned} \phi_z * \frac{\chi_1}{2k} &= \frac{P_1(z)}{e_1} + \sum_{n=1}^{\infty} \left(\frac{P_{n-1}(z)}{2ke_{n-1}} + \frac{\omega^2 P_{n+1}(z)}{2ke_{n+1}} \right) \chi_n \\ &= \frac{z}{2k} \sum_{n=0}^{\infty} \frac{P_n(z)}{e_n} \chi_n \\ &= \frac{z}{2k} \phi_z. \end{aligned}$$

By [4, Corollary 1.7] ϕ_z is spherical, equivalently it is a multiplicative linear functional on $(\mathbb{C}F_k)_r$. We can now prove

Proposition 4.2. *Let* $\alpha \in (\mathbb{C}F_k)_r$ *and let* $z \in \mathbb{C}$. *Then* $\alpha * \phi_z = 0$ *if and only if* $\langle \chi_n * \alpha, \overline{\phi_z} \rangle = 0$ *for all integers* $n \geq 0$.

Proof. Using $\chi_n * \alpha = \sum_{g \in E_n} L_g \alpha$ with Lemma 2.2, $\langle \chi_n * \alpha, \overline{\phi_z} \rangle = 0$ for each $n \geq 0$.

Conversely, write $\alpha = \sum_{j=0}^{m} a_j \chi_j$. Then

$$\begin{aligned} \langle \alpha, \overline{\phi_z} \rangle &= \sum_{j,n} \frac{a_j P_n(z)}{e_n} \langle \chi_j, \chi_n \rangle \\ &= \sum_{j=0}^{m} a_j P_j(z) = \hat{\alpha}(z). \end{aligned}$$

The function $\overline{\phi_z}$ is multiplicative on $(\mathbb{C}F_k)_r$ because $\overline{\phi_z} = \phi_{\overline{z}}$. Thus $\langle \chi_n * \alpha, \overline{\phi_z} \rangle = \hat{\alpha}(z) P_n(z)$. Lemma 4.1 allows us to deduce that if $x \in F_k$ and $|x| = n$, then $\langle x * \alpha, \overline{\phi_z} \rangle = \hat{\alpha}(z) \frac{P_n(z)}{e_n}$. The result now follows from Lemma 2.2. □

Corollary 4.3. *Let* $0 \neq \alpha \in (\mathbb{C}F_k)_r$ *and let* $z \in \mathbb{C}$. *Then* $\alpha * \phi_z = 0$ *if and only if* $z \in Z(\alpha)$.

Proof. We saw in the proof of the above proposition that $\langle \chi_n * \alpha, \overline{\phi_z} \rangle = \hat{\alpha}(z) P_n(z)$ for each positive integer n. The corollary now follows. □

We need the following fact about maximal ideals in $(\mathbb{C}F_k)_r$.

Proposition 4.4. *If M is a maximal ideal in $(\mathbb{C}F_k)_r$, then $(\mathbb{C}F_k)_r/M$ is isomorphic to $\mathbb{C}$.*

Proof. Let $\mathbb{C}[x]$ be the polynomial ring over $\mathbb{C}$. The proposition will be proved if we can show $(\mathbb{C}F_k)_r$ is isomorphic to $\mathbb{C}[x]$, since $\mathbb{C}[x]/J$ is isomorphic to $\mathbb{C}$ for each maximal ideal J in $\mathbb{C}[x]$. For $\alpha = \sum_{i=0}^n a_i \chi_i \in (\mathbb{C}F_k)_r$, define $T\alpha \in \mathbb{C}[x]$ by

$$T\alpha = \sum_{i=0}^n a_i p_i(x).$$

We will now show that T gives the desired isomorphism. Clearly T is linear. Note that if $z \in \mathbb{C}$, then $T\alpha(z) = \langle \phi_z, \alpha \rangle$. Combining this observation with the fact ϕ_z is multiplicative on $(\mathbb{C}F_k)_r$ for each $z \in \mathbb{C}$, we obtain $T(\alpha * \beta) = (T\alpha)(T\beta)$, where $\beta \in (\mathbb{C}F_k)_r$. If $T\alpha = \sum_{i=0}^n a_i p_i(x) = 0$, then it must be the case $a_0 = a_1 = \cdots = a_n = 0$. Thus T is one-to-one. Let $p \in \mathbb{C}[x]$. We may assume without loss of generality that p is a monic polynomial. Write $p = (x - c_1)(x - c_2) \cdots (x - c_n)$, where $c_k \in \mathbb{C}$ and n is the degree of p. The function $(\chi_1 - c_1) * (\chi_1 - c_2) * \cdots * (\chi_1 - c_n) \in (\mathbb{C}F_k)_r$ and T maps this element to p, so T is onto. Hence T is an isomorphism between $(\mathbb{C}F_k)_r$ and $\mathbb{C}[x]$ and the proposition now follows □

We can now state and prove the following Wiener type result for F_k, which corresponds to Theorem 3.2.

Theorem 4.5. *Let F_k be the free group with $2 \leq k \leq \mathbb{N}$. Let I be an ideal in $(\mathbb{C}F_k)_r$. Then $Z(I) = \emptyset$ if and only if $I = (\mathbb{C}F_k)_r$.*

Proof. Suppose $Z(I) \neq \emptyset$ and let $z \in Z(I)$. Then for each $\alpha \in I, \alpha * \phi_z = 0$ by Corollary 4.3. Since $\chi_0 * \phi_z \neq 0$ it follows that $I \neq (\mathbb{C}F_k)_r$.

Conversely, assume $I \neq (\mathbb{C}F_k)_r$. Let M be a maximal ideal in $(\mathbb{C}F_k)_r$ that contains I. Then $(\mathbb{C}F_k)_r/M$ is isomorphic to $\mathbb{C}$. Hence, there is a ring homomorphism $\psi \colon (\mathbb{C}F_k)_r \mapsto \mathbb{C}$ for which the kernel of ψ is M. The multiplicative linear functional ψ is determined by its value on χ_1. Set $z = \psi(\chi_1)$. Because $\chi_n = P_n(\chi_1)$, we obtain $\psi(\chi_n) = P_n(\psi(\chi_1)) = P_n(z)$. Consequently, $\psi = \phi_z$. Furthermore, if $\alpha \in I$, then $\langle \chi_n * \alpha, \overline{\phi_z} \rangle = 0$ for each nonnegative integer n. By Proposition 4.2 and Corollary 4.3, $z \in Z(\alpha)$ for each $\alpha \in I$. Therefore, $z \in Z(I)$. □

The following result, which is similar to Corollary 3.3, gives necessary and sufficient conditions for a collection of radial subsets of F_k to have the Pompeiu property.

Corollary 4.6. *Let $2 \leq k \in \mathbb{N}$ and let $\mathcal{K}$ be a finite set of nonempty finite radial subsets of F_k. The $\mathcal{K}$ has the Pompeiu property if and only if $\cap_{K \in \mathcal{K}} Z(\chi_K) = \emptyset$.*

Proof. First assume $\cap_{K \in \mathcal{K}} Z(\chi_K) \neq \emptyset$ and let $z \in \cap_{K \in \mathcal{K}} Z(\chi_K)$. Then $\chi_K * \phi_z = 0$ for each $K \in \mathcal{K}$. Since χ_K and ϕ_z are both radial, it follows from Proposition 2.3 that $\mathcal{K}$ does not have the Pompeiu property.

Now assume $\cap_{K \in \mathcal{K}} Z(\chi_K) = \emptyset$. Let I be the ideal in $(\mathbb{C}F_k)_r$ generated by $\{\chi_K\}_{K \in \mathcal{K}}$. Then $Z(I) = \emptyset$ and by Theorem 4.5 $I = (\mathbb{C}F_k)_r$. If $f \in \mathcal{F}(F_k)$ and $\chi_K * f = 0$ for each $K \in \mathcal{K}$, then $\chi_0 * f = 0$. Hence, it must the case $f = 0$ and $\mathcal{K}$ has the Pompeiu property. □

A special case of the Pompeiu problem is the two circle problem. Before we state this problem we will define some needed terms. For the rest of this paragraph we will assume the G is a finitely generated group with generating set $x_1, \dots, x_n$. The Cayley graph Γ of G is the graph whose vertices are the elements of G, and $g, h \in G$ are connected by an edge if and only if $h = gx_i^{\pm 1}$ for some generator x_i of G. Note that Γ is a connected graph. Now G is a metric space via the shortest path metric on Γ. That is, if $x, y \in G$, then $d(x, y)$ equals the number of edges of the shortest path in Γ joining x and y. We now state the two circle problem for discrete groups.

Problem 4.7. *Let r and s be natural numbers. Is $f = 0$ the only function on G that satisfies*

$$\sum_{d(x,y)=r} f(y) = 0 = \sum_{d(x,y)=s} f(y) \tag{4.1}$$

for all $x \in G$?

Cohen and Picardello studied this problem in [3] for homogeneous trees. The Cayley graph of the free group $F_k, 2 \leq k \in \mathbb{N}$, is a homogeneous tree of degree $2k$. We now proceed to explain how our results overlap with some of the results in [3].

Let K_n be the set of elements in F_k that have length n. Each element of K_n has distance n from the identity element e of F_k. From a geometric point of view we can think of K_n as a sphere in F_k of radius n with center e. Denote the characteristic function on K_n by χ_n. Set

$$\phi_0 = \sum_{n=0}^{\infty} \frac{P_n(0)}{e_n} \chi_n = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2k-1)^n} \chi_{2n}.$$

Since $\{0\} = Z(\chi_1)$, Corollary 4.3 says that $\chi_1 * \phi_0 = 0$. Furthermore, $\phi_0 \in L^p(F_k)$ for $p > 2$. Because χ_1 and ϕ_0 are both radial, $\chi_1 * \widetilde{\phi_0} = \chi_1 * \phi_0 = 0$. Thus ϕ_0 satisfies (1.1) by Proposition 2.3. So K_1 does not have the Pompeiu property when the class of functions under consideration contains $L^p(F_k)$ for some $p > 2$. In fact, if n an odd natural number, then $0 \in Z(\chi_n)$ due to $\widehat{\chi_n} = P_n(z)$ being an odd polynomial. Thus $\chi_n * \widetilde{\phi_0} = 0$ and K_n does not have the Pompeiu property for all odd natural numbers. We now move on to the two circle problem.

Let r and s be natural numbers and set $\mathcal{K} = \{K_r, K_s\}$. Observe that if $x \in F_k$, then xK_r and xK_s are spheres in F_k with center x and radius r and s respectively. Consequently, $f = 0$ is the only solution to (4.1) if and only if $\mathcal{K}$ has the Pompeiu property. Corollary 4.6 tells us $\mathcal{K}$ has the Pompeiu property precisely when $\widehat{\chi_s} = P_s(z)$ and $\widehat{\chi_r} = P_r(z)$ do not have a common root. If r and s are both odd, then as we saw in the previous paragraph 0 is a common root of both $\widehat{\chi_s}$ and $\widehat{\chi_r}$. Thus $\mathcal{K}$ does not have the Pompeiu property and ϕ_0 is a nonzero function that satisfies (4.1). This is essentially Theorem 1(a) of [3] for the case of homogeneous trees with positive even degree.

Now suppose r and s are both not odd. It was shown in [3, Section 2] that $\widehat{\chi_r} = P_r(z)$ and $\widehat{\chi_s} = P_s(z)$ do not have a common root. Thus $\mathcal{K}$ has the Pompeiu property and $f = 0$ is the only solution to (4.1).

References

[1] Carlos A. Berenstein and Lawrence Zalcman. Pompeiu's problem on symmetric spaces. *Comment. Math. Helv.*, 55(4):593–621, 1980.

[2] Joel M. Cohen. von Neumann dimension and the homology of covering spaces. *Quart. J. Math. Oxford Ser. (2)*, 30(118):133–142, 1979.

[3] Joel M. Cohen and Massimo A. Picardello. The 2-circle and 2-disk problems on trees. *Israel J. Math.*, 64(1):73–86, 1988.

[4] Alessandro Figà-Talamanca and Massimo A. Picardello. *Harmonic analysis on free groups*, volume 87 of *Lecture Notes in Pure and Applied Mathematics*. Marcel Dekker Inc., New York, 1983.

[5] M. Laczkovich and G. Székelyhidi. Harmonic analysis on discrete abelian groups. *Proc. Amer. Math. Soc.*, 133(6):1581–1586 (electronic), 2005.

[6] P. A. Linnell. Zero divisors and group von Neumann algebras. *Pacific J. Math.*, 149(2):349–363, 1991.

[7] Peter A. Linnell. Zero divisors and $L^2(G)$. *C. R. Acad. Sci. Paris Sér. I Math.*, 315(1):49–53, 1992.

[8] Peter A. Linnell and Michael J. Puls. Zero divisors and $L^p(G)$. II. *New York J. Math.*, 7:49–58 (electronic), 2001.

[9] Norbert Peyerimhoff and Evangelia Samiou. Spherical spectral synthesis and two-radius theorems on Damek-Ricci spaces. *Ark. Mat.*, 48(1):131–147, 2010.

[10] Michael J. Puls. Zero divisors and $L^p(G)$. *Proc. Amer. Math. Soc.*, 126(3):721–728, 1998.

[11] Rama Rawat and Alladi Sitaram. The injectivity of the Pompeiu transform and L^p-analogues of the Wiener-Tauberian theorem. *Israel J. Math.*, 91(1-3):307–316, 1995.

[12] David Scott and Alladi Sitaram. Some remarks on the Pompeiu problem for groups. *Proc. Amer. Math. Soc.*, 104(4):1261–1266, 1988.

[13] Lawrence Zalcman. Offbeat integral geometry. *Amer. Math. Monthly*, 87(3):161–175, 1980.

Department of Mathematics, John Jay College-CUNY, 445 West 59th Street, New York, NY 10019, USA

E-mail address: `mpuls@jjay.cuny.edu`